STROBE Statement—Checklist of items that should be included in reports of *cross-sectional studies* 

	Item No	Recommendation
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract
Title and abstract	-	[Abstract – methods, p.2]
		(b) Provide in the abstract an informative and balanced summary of what was done
		and what was found [Abstract- methods and results, p.2]
Introduction		
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported
		[p.3]
Objectives	3	State specific objectives, including any prespecified hypotheses [p.3-4]
Methods		
Study design	4	Present key elements of study design early in the paper [Methods- study
		population, p.4; statistical analysis – p.6]
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment,
		exposure, follow-up, and data collection [Methods- study population, p.4]
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of
		participants [Methods- study population, p.4]
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect
		modifiers. Give diagnostic criteria, if applicable [Methods- definitions, p.4-6]
Data sources/	8*	For each variable of interest, give sources of data and details of methods of
measurement		assessment (measurement). Describe comparability of assessment methods if there is
		more than one group [Methods- definitions, p.4-5]
Bias	9	Describe any efforts to address potential sources of bias
Study size	10	Explain how the study size was arrived at [Abstract- p.2; Methods- study
		population, p.4; Table 1]
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable,
		describe which groupings were chosen and why [Methods- definitions, p.4-6]
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding
		[Methods- statistical analysis, p.6-7]
		(b) Describe any methods used to examine subgroups and interactions [Methods-
		statistical analysis, p.6-7]
		(c) Explain how missing data were addressed [N/A]
		(d) If applicable, describe analytical methods taking account of sampling strategy
		[Methods- statistical analysis, p.6]
		$(\underline{e})$ Describe any sensitivity analyses [N/A]
Results		
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially
		eligible, examined for eligibility, confirmed eligible, included in the study,
		completing follow-up, and analysed [Abstract, p.2, Table 1]
		(b) Give reasons for non-participation at each stage [N/A]
		(c) Consider use of a flow diagram [N/A]
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and
		information on exposures and potential confounders [Results, p.7]
		(b) Indicate number of participants with missing data for each variable of interest
		[N/A]
Outcome data	15*	Report numbers of outcome events or summary measures [Results- Trends in

l estimates and, if applicable, confounder-adjusted estimates and
95% confidence interval). Make clear which confounders were
ny they were included [Results-, p.7-9]
boundaries when continuous variables were categorized [N/A]
ider translating estimates of relative risk into absolute risk for a
riod [N/A]
es done—eg analyses of subgroups and interactions, and
[Results- p.7-9]
ults with reference to study objectives [Interpretation- Main
of the study, taking into account sources of potential bias or
ss both direction and magnitude of any potential bias
imitations, p.12]
erall interpretation of results considering objectives, limitations,
yses, results from similar studies, and other relevant evidence
10-11]
isability (external validity) of the study results [Interpretation,
funding and the role of the funders for the present study and, if
original study on which the present article is based [p.1]

<sup>\*</sup>Give information separately for exposed and unexposed groups.

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at http://www.plosmedicine.org/, Annals of Internal Medicine at http://www.annals.org/, and Epidemiology at http://www.epidem.com/). Information on the STROBE Initiative is available at www.strobe-statement.org.